

RESTRICTED 1-3-2 PERMUTATIONS AND GENERALIZED PATTERNS

TOUFIK MANSOUR

LaBri, Université Bordeaux I,
351 cours de la Libération, 33405 Talence Cedex
toufik@labri.fr

ABSTRACT

Recently, Babson and Steingrímsson (see [BS]) introduced generalized permutations patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation.

We study generating functions for the number of permutations on n letters avoiding 1-3-2 (or containing 1-3-2 exactly once) and an arbitrary generalized pattern τ on k letters, or containing τ exactly once. In several cases the generating function depends only on k and is expressed via Chebyshev polynomials of the second kind, and generating function of Motzkin numbers.

1. INTRODUCTION

Permutation patterns: let $\alpha \in S_n$ and $\tau \in S_k$ be two permutations. We say that α *contains* τ if there exists a subsequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $(\alpha_{i_1}, \dots, \alpha_{i_k})$ is order-isomorphic to τ ; in such a context τ is usually called a *pattern*. We say α *avoids* τ , or is τ -*avoiding*, if such a subsequence does not exist. The set of all τ -avoiding permutations in S_n is denoted $S_n(\tau)$. For an arbitrary finite collection of patterns T , we say that α avoids T if α avoids any $\tau \in T$; the corresponding subset of S_n is denoted $S_n(T)$.

While the case of permutations avoiding a single pattern has attracted much attention, the case of multiple pattern avoidance remains less investigated. In particular, it is natural, as the next step, to consider permutations avoiding pairs of patterns τ_1, τ_2 . This problem was solved completely for $\tau_1, \tau_2 \in S_3$ (see [SS]), for $\tau_1 \in S_3$ and $\tau_2 \in S_4$ (see [W]), and for $\tau_1, \tau_2 \in S_4$ (see [B, K] and references therein). Several recent papers [CW, MV1, Kr, MV2, MV3] deal with the case $\tau_1 \in S_3, \tau_2 \in S_k$ for various pairs τ_1, τ_2 . Another natural question is to study permutations avoiding τ_1 and containing τ_2 exactly t times. Such a problem for certain $\tau_1, \tau_2 \in S_3$ and $t = 1$ was investigated in [R], and for certain $\tau_1 \in S_3, \tau_2 \in S_k$ in [RWZ, MV1, Kr, MV2]. The tools involved in these papers include continued fractions, Chebyshev polynomials, and Dyck paths.

Generalized permutation patterns: in [BS] Babson and Steingrímsson introduced generalized permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. The idea for

Babson and Steingrímsson in introducing these patterns was study of Mahonian statistics, and they proved that all Mahonian permutation statistics in the literature can be written as linear combinations of such patterns.

We define our *generalized patterns* as words in the alphabet $1, 2, 3, \dots$, where two adjacent letters may or may not be separated by a dash. The absence of a dash between two adjacent letters in a pattern indicates that the corresponding letters in the permutation must be adjacent, and in the order (order-isomorphic) given by the pattern. An example of a generalized pattern is 1-23. An 1-23 subword of a permutation $\pi = (\pi_1, \pi_2, \dots, \pi_n)$ is a subword $(\pi_i, \pi_j, \pi_{j+1})$ ($i < j$), such that $\pi_i < \pi_j < \pi_{j+1}$.

Claesson [C] presented a complete solution for the number of permutations avoiding any single generalized pattern of length three with exactly one adjacent pair of letters, and presented some results for the number of permutations avoiding two patterns of length three with exactly one adjacent pair of letters. Claesson and Mansour [CM] presented a complete solution for the number of permutations avoiding any double generalized patterns of length three with exactly one adjacent pair of letters. Besides, Kitaev [Ki] investigate simultaneous avoidance of two or more 3-letters generalized patterns without internal dashes.

A *Motzkin path* of length n is a lattice path from $(0, 0)$ to $(n, 0)$ with steps $(1, 0)$, $(1, 1)$, and $(1, -1)$ that never goes below the a -axis. Letting L , U , and D represent the steps $(1, 0)$, $(1, 1)$, and $(1, -1)$ respectively, we code such a path with word over $\{L, U, D\}$. If p any nonempty Motzkin path, then p can be decomposed as either, $p = Lp'$ or $p = Up'Dq'$ where p', q' are Motzkin paths. The n th Motzkin number M_n is the number of Motzkin paths of length n . The first few of the M_n are

$$1, 1, 2, 4, 9, 21, 51, 127, 323, 835, 2188, 5798, 15511, 41835, 113634, 310572.$$

Notation 1.1. The generating function for Motzkin numbers M_n we denote by $M(x)$; that is, $M(x) = \sum_{n \geq 0} M_n x^n$.

A *Chebyshev polynomials of the second kind* are defined $U_p(\cos \theta) = \sin(p + 1)\theta / \sin \theta$. Following [CW, MV1, Kr, MV2, MV3], introduce

$$R_p(x) = \frac{U_{p-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_p\left(\frac{1}{2\sqrt{x}}\right)}.$$

As consequence to [MV2], we present a general approach to the study of permutations avoiding 1-3-2 and avoiding a generalized pattern in S_k (or containing exactly once). As a consequence, we derive many of the previously known results for this kind of problems, as well as many new results.

The paper is organized as follows. The case of permutations avoiding both 1-3-2 and τ generalized pattern is treated in Section 2. We derive a simple recursion for the corresponding generating function for general τ . This recursion can be solved explicitly for several interesting cases. The case of permutations avoiding 1-3-2 and containing τ once is treated in Section 3. Here again we start from general recursion, and then solve it for several particular cases. The case containing 132 exactly once is treated in section 5 and 6 for avoiding τ , or containing τ exactly once; respectively.

Most of the explicit solutions obtained in Sections 2 and 3 involve Chebyshev polynomials of the second kind and generating function of Motzkin numbers.

2. AVOIDING 1-3-2 AND A GENERALIZED PATTERN

Consider an arbitrary pattern $\phi = (\phi_1, \dots, \phi_k) \in S_k(1-3-2)$. Recall that ϕ_i is said to be a *right-to-left maximum* if $\phi_i > \phi_j$ for any $j > i$. Let $m_0 = k, m_1, \dots, m_r$ be the right-to-left maxima of ϕ written from left to right. Then ϕ can be represented as

$$\phi = (\phi^0, m_0, \phi^1, m_1, \dots, \phi^r, m_r),$$

where each of ϕ^i may be eventually empty, and all the entries of ϕ^i are greater than all the entries of ϕ^{i+1} . This representation is called the *canonical decomposition* of ϕ . Given the canonical decomposition, we define the i th *prefix* of ϕ by $\pi^i = (\phi^0, m_0, \dots, \phi^i, m_i)$ for $1 \leq i \leq r$ and $\pi^0 = \phi^0, \pi^{-1} = \emptyset$. Besides, the i th *suffix* of ϕ is defined by $\sigma^i = (\phi^i, m_i, \dots, \phi^r, m_r)$ for $0 \leq i \leq r$ and $\sigma^{r+1} = \emptyset$. The following representation $\tau = \phi^0 - m_0 - \phi^1 - m_1 - \dots - \phi^r - m_r$ is called the *canonical decomposition for generalized pattern* $\phi \in S_k(1-3-2)$. Similarly we define π^i and σ^i for τ .

Let $f_\tau(n)$ denote the number of permutations in $S_n(1-3-2)$ such that avoiding τ , and let $F_\tau(x) = \sum_{n \geq 0} f_\tau(n)x^n$ be the corresponding generating function. By $f_\tau^\beta(n)$ we denote the number of permutations in $S_n(1-3-2)$ avoiding τ and containing β . The following proposition is the base of all the other results in this Section, which is obtained by use the same proof argument of [MV2, Th. 1].

Theorem 2.1. *Let $\phi \in S_k(1-3-2)$ such that $\phi = (\phi^0, m_0, \phi^1, m_1, \dots, \phi^r, m_r)$, and let $\tau = (\tau^0 - m_0 - \tau^1 - m_1 - \dots - \tau^r - m_r)$ where $\phi^i = \overline{\tau^i}$. Then*

$$F_\tau(x) = 1 + x \sum_{j=0}^r (F_{\pi^j}(x) - F_{\pi^{j-1}}(x)) F_{\sigma^j}(x).$$

Example 2.2. (Cleasson and Mansour [CM]) *Let us find $F_{12-3}(x)$. The canonical decomposition for 12-3 is (12)-3, so $r = 0$, and hence $F_{12-3}(x) = 1 + xF_{12}(x)F_{12-3}(x)$. By definitions, $F_{12}(x) = \frac{1}{1-x}$, hence $F_{12-3}(x) = \frac{1-x}{1-2x}$. similarly, since $F_{21}(x) = \frac{1}{1-x}$ we get that $F_{21-3}(x) = \frac{1-x}{1-2x}$.*

The case of varying k is more interesting. As an extension of Example 2.2.

Theorem 2.3. *Let $\tau \in \{12, 1-2, 21, 2-1\}$; for any $k \geq 2$,*

$$F_{\tau-3-4-\dots-k}(x) = R_k(x).$$

Proof. By Theorem 2.1 it is easy to see

$$F_{\tau-3-\dots-k}(x) = \frac{1}{1 - xF_{\tau-3-\dots-(k-1)}(x)}.$$

Besides, by definitions $F_\tau(x) = \frac{1}{1-x}$, hence by use [MV2, Eq. 4.1] the theorem holds. $\square$

Notation 2.4. *The generalized pattern $123 \dots k$ we denote by $[k]$.*

Theorem 2.5. *The generating function $F_{[k]}(x)$ is defined by the following equation*

$$F_{[k]}(x) = \sum_{j=0}^{k-1} (xF_{[k]}(x))^j.$$

Proof. Let $\alpha \in S_n(1-3-2, [k])$; If α nonempty, then we may write $\alpha = \beta^1 n \gamma$ such that $\alpha_a = n$ where β^1 and γ avoids 1-3-2, β^1 is a permutation of $\{n-1, n-2, \dots, n-a+1\}$, and γ is a permutation of $\{1, 2, \dots, n-a\}$. If β^1 is nonempty, then let us assume that $\beta_{a-1}^1 = m_1$ and since β^1 avoids 1-3-2, we may write $\beta^1 = \delta^1 \beta^2 m_1$ where β^2 and δ^1 avoids 1-3-2, δ^1 is a permutation of $\{n-1, \dots, m_1+1\}$, and β^2 is a permutation of $\{m_1-1, \dots, n-a+1\}$, and so on β^2 . Thus we may write $\beta^1 = \delta^1 \delta^2 \dots \delta^{k-2}, m_{k-2}, \dots, m_1$ where δ^j avoids 1-3-2 and $[k]$ for any $j \leq k-2$, and $\delta^{k-1} = \emptyset$. Therefore, in terms of generating function it is easy to get that

$$F_{[k]}(x) = 1 + xF_{[k]}(x) + \dots + x^{k-1}F_{[k]}^{k-1}(x).$$

□

Example 2.6. (see Barcucci, Del Lungo, Pergola, and Pinzani [BDPP]) *Theorem 2.5 yields $F_{[1]}(x) = 1$, and $F_{[2]}(x) = \frac{1}{1-x}$. The first interesting case which Theorem 2.5 yields $F_{123}(x) = M(x)$. Also by use Theorem 2.1 we have*

$$F_{123-4}(x) = \frac{1}{1 - xF_{123}(x)} = \frac{1}{2} \sqrt{\frac{1+x}{1-3x}},$$

which is the generating function of Directed animals of size n .

Similarly, by proof of Theorem 2.5 we get the following.

Theorem 2.7. *Let $k > d \geq 1$ and $\tau = \tau' - (d+1)(d+2) \dots k$. Then*

$$F_\tau(x) = \sum_{j=0}^{k-d-1} (xF_\tau(x))^j + x^{k-d} F_\tau^{k-d}(x) F_{\tau'}(x).$$

Example 2.8. (Claesson [C, Pro. 16]) *By Theorem 2.7 we obtained that $F_{1-23}(x) = 1 + xF_{1-23}(x) + x^2 F_{1-23}(x)$, hence $F_{1-23}(x) = M(x)$, the generating function of Motzkin numbers.*

The case of varying k is more interesting. As an extension of Example 2.8.

Theorem 2.9. *For all $k \geq 3$, the generating functions $F_{1-2-\dots-(k-2)-(k-1)k}(x)$, $F_{12-3-\dots-(k-2)-(k-1)k}(x)$, $F_{2-1-3-\dots-(k-2)-(k-1)k}(x)$, $F_{21-3-\dots-(k-2)-(k-1)k}(x)$ are given by*

$$\frac{1 - x - \sqrt{1 - 2x + x^2 - 4x^2 R_{k-2}(x)}}{2x^2 R_{k-2}(x)}.$$

Proof. By Theorem 2.7

$$F_{1-2-\dots-(k-2)-(k-1)k}(x) = 1 + xF_{1-2-\dots-(k-2)-(k-1)k}(x) + x^2 F_{1-2-\dots-(k-2)-(k-1)k}^2(x) F_{1-2-\dots-(k-2)}(x).$$

Therefore, by use Theorem 2.3 the theorem holds in this case. Similarly for $12-3-\dots-(k-2)-(k-1)k$, for $2-1-3-\dots-(k-2)-(k-1)k$, and for $21-3-\dots-(k-2)-(k-1)k$. □

Example 2.10. *Theorem 2.9 yields that, the generating functions $F_{12-34}(x)$, $F_{21-34}(x)$, $F_{1-2-34}(x)$, and $F_{2-1-34}(x)$ are given by*

$$\frac{1 - 2x + x^2 - \sqrt{1 - 4x + 2x^2 + x^4}}{2x^2}.$$

Notation 2.11. The generalized pattern $k \dots 321$ we denote by $\langle k \rangle$.

So, by the same way of proof Theorem 2.5 we get the following.

Theorem 2.12. The generating function $F_{\langle k \rangle}(x)$ is defined by the following equation

$$F_{\langle k \rangle}(x) = \sum_{j=0}^{k-1} (xF_{\langle k \rangle}(x))^j.$$

Proof. Let $\alpha \in S_n(1-3-2, \langle k \rangle)$; If α nonempty, then we may write $\alpha = \gamma n \beta^1$ such that $\alpha_a = n$ where β^1 and γ avoids 1-3-2, γ is a permutation of $\{n-1, n-2, \dots, n-a+1\}$, and β^1 is a permutation of $\{1, 2, \dots, n-a\}$. If β^1 is nonempty, then let us assume that $\beta_1^1 = m_1$ and since β^1 avoids 1-3-2, we may write $\beta^1 = m_1 \beta^2 \delta^1$ where β^2 and δ^1 avoids 1-3-2, δ^1 is a permutation of $\{n-a, \dots, m_1+1\}$, and β^2 is a permutation of $\{m_1-1, \dots, 1\}$, and so on β^2 . Thus we may write $\beta^1 = m_1, \dots, m_{k-2}, \delta^{k-1}, \delta^{k-2} \delta^{k-3} \dots \delta^1$ where δ^j avoids 1-3-2 and $\langle k \rangle$ for any $j \leq k-2$, and since α avoids $\langle k \rangle$, then $\delta^{k-1} = \emptyset$. Therefore, in terms of generating function it is easy to get that

$$F_{[k]}(x) = 1 + xF_{[k]}(x) + \dots + x^{k-1}F_{[k]}^{k-1}(x).$$

□

Example 2.13. By Theorem 2.12 we get $F_{21}(x) = \frac{1}{1-x}$. The first interesting special case for $k = 3$,

$$F_{321}(x) = 1 + xF_{321}(x) + x^2F_{321}^2(x),$$

hence $F_{321}(x) = M(x)$. Besides, by use Theorem 2.1 we have

$$F_{321-4}(x) = \frac{1}{1 - xF_{321}(x)} = \frac{1}{2} \sqrt{\frac{1+x}{1-3x}},$$

which is the generating function of Directed animals of size n (see [BDPP]).

Another way to generalize these result by looking at patterns of the form $(d+1, \dots, k, 1, 2, \dots, d) \in S_k$.

Example 2.14. By Theorem 2.1 we get

$$F_{45-6-12-3}(x) = 1 + xF_{21}(x)F_{45-6-12-3}(x) + x(F_{45-6-12-3}(x) - F_{12}(x))F_{12-3}(x),$$

which means that

$$F_{45-6-12-3}(x) = \frac{(1-3x)(1-x)}{1-5x+6x^2-x^3}.$$

This example can be generalized by use Theorem 2.1 as following.

Theorem 2.15. Let $k > d$, and let $\tau = \tau' - k - \tau'' - d$ be generalized pattern such that the minimal letter of τ' is $d+1$. Then

$$F_{\tau}(x) = \frac{1}{1 - x \frac{1 - xF_{\tau'}(x)F_{\tau''}(x)}{1 - x(F_{\tau'}(x) + F_{\tau''}(x))}}.$$

Example 2.16. Let $\tau = (d+1)(d+2)-(d+3)-\dots-k-12-3-\dots-d$ where $d \geq 2, k \geq 4$. By Theorem 2.15 with use 2.3

$$F_\tau(x) = \frac{1}{1 - x \frac{1-xR_{k-d}(x)R_d(x)}{1-x(R_{k-d}(x)+R_d(x))}},$$

and [MV2, Section 4] yields $F_\tau(x) = R_k(x)$.

For a further generalization of the above example, consider the following definition. We say that τ is a *generalized wedge pattern* if it can be represented as $\tau = (\phi^1, \psi^1, \dots, \phi^r, \psi^r)$ so that $\bar{\tau}$ is a *wedge pattern* (see [MV2, ?]) and each ϕ^j and ψ^j is either $a-(a+1)-\dots-b$ or $a(a+1)-(a+2)-\dots-b$. For example, 6-4-5-7-8-3-9-12 and 45-6-3-7-8-12-9 are generalized wedge patterns. The following theorem is analogue of Mansour and Vainshtein result [MV3, Th. 2.6]

Theorem 2.17. $F_\tau(x) = R_k(x)$ for any generalized wedge pattern τ of k -letters.

Remark 2.18. A comparison of Theorem 2.17 with the Main result of [MV3] suggests that there should exist a bijection between the sets $S_n(3-2-1, (m+1)-\dots-k-1-\dots-m)$ and $S_n(1-3-2, \tau)$ for any generalized wedge pattern τ . However, we failed to produce such a bijection, and finding it remains a challenging open question.

Corollary 2.19. 1. $F_{123}(x) = F_{321}(x) = \frac{1-x-\sqrt{1-2x-3x^2}}{2x^2}$;
 2. $F_{132}(x) = \frac{1-\sqrt{1-4x}}{2x}$;
 3. $F_{213}(x) = F_{312}(x) = \frac{1-x^2-\sqrt{(1+x^2)^2-4x}}{2x(1-x)}$;
 4. $F_{231}(x) = \frac{1-x}{1-2x}$.

Proof. The case (1) holds by Theorem 2.5, and Theorem 2.12; the case (2) by definition and Knuths results [Kn].

To verify (3); let $\alpha \in S_n(213, 1-3-2)$; we may write $\alpha = \beta n \gamma$ such that $\alpha_k = n$, where β and γ are avoids 1-3-2, β is a permutation of $\{n-k+1, \dots, n-2, n-1\}$, and γ is a permutation of $\{1, 2, \dots, n-k\}$. Let m the maximal such that the last m numbers in β are increasing. So, since β avoids 1-3-2 we obtain $\beta = (\delta^1, \dots, \delta^m, a^m, \dots, a^1)$ where $a^{j+1} < a^j$ and every element in δ^j is bigger that every element in δ^{j+1} . On the other hand, (β, n) avoids 213, so $\delta^m = \emptyset$. Hence

$$F_{213}(x) = 1 + xF_{213}(x) + x^2F_{213}(x) + x^3F_{213}^2(x) + x^4F_{213}^3(x) + \dots,$$

which means that

$$F_{213}(x) = 1 + xF_{213}(x) + \frac{x^2F_{213}(x)}{1 - xF_{213}(x)}.$$

Similarly, we obtain $F_{312}(x) = F_{213}(x)$.

Finally, by use the same argument above proof we easy to get $F_{231}(x) = 1 + x(F_{231}(x) - 1) + xF_{231}(x)$, which means that $F_{231}(x) = \frac{1-x}{1-2x}$. $\square$

3. AVOIDING 1-3-2 AND CONTAINING ONCE A GENERALIZED PATTERN

Let $g_\tau(n)$ denote the number of permutations in $S_n(1-3-2)$ that contains τ exactly once where τ generalized pattern avoids 1-3-2, and $g_\tau^\phi(n)$ denote the number of permutations in $S_n(1-3-2, \phi)$ that contains τ exactly once. We denote by $G_\tau(x)$ and $G_\tau^\phi(x)$ the corresponding ordinary generating functions.

The following statement is the analog of [MV2, Th. 3.1], which is yields immediately by the same argument proof of [MV2, Th. 3.1].

Theorem 3.1. Let $\phi \in S_k(1-3-2)$ such that the canonical decomposition of ϕ given by $\phi = (\phi^0, m_0, \phi^1, m_1, \dots, \phi^r, m_r)$, and let $\tau = (\theta^0 - m_0 - \theta^1 - m_1 - \dots - \theta^r - m_r)$ where $\phi^i = \bar{\theta}^i$. Then

$$(1 - xF_{\pi^0}(x) - xF_{\sigma^r}(x))G_\tau(x) = x \sum_{j=1}^r G_{\pi^{j-1}}^{\pi^j}(x) G_{\sigma^j}^{\sigma^{j-1}}(x)$$

for $r \geq 1$, and

$$G_\tau(x) = \frac{x F_\tau(x) G_{\pi^0}(x)}{1 - x F_{\pi^0}(x)}$$

for $r = 0$.

Theorem 3.2. Let $k \geq 1$,

$$G_{[k]}(x) = \sum_{j=1}^{k-1} j x^j G_{[k]}(x) F_{[k]}^{j-1}(x) + x^k F_{[k]}^k(x),$$

where $F_{[k]}(x)$ given in Theorem 2.5.

Proof. By use the argument proof of Theorem 2.5 and Theorem 2.12, the rest is easy to see. $\square$

Example 3.3. Theorem 3.2, and Theorem 2.3 yields $G_{12}(x) = G_{21}(x) = \frac{x^2}{(1-x)^3}$. Theorem 3.2, Example 2.6 and Example 2.13 yields

$$G_{123}(x) = G_{321}(x) = \frac{x^3 M^3(x)}{1 - x - 2x^2 M(x)} = 2x - 1 + \frac{1 - 3x + 2x^3}{2x^3 \sqrt{1 - 2x - 3x^2}}.$$

Corollary 3.4. For any $k \geq 2$;

$$G_{12-3-\dots-k}(x) = \frac{1}{(1-x)U_k^2\left(\frac{1}{2\sqrt{x}}\right)}.$$

Proof. Let $\tau = 12-3-\dots-k$; then $r = 0$, and it follows from Theorem 3.1 that

$$G_{12-3-\dots-k}(x) = \frac{x F_{12-3-\dots-k}(x) G_{12-3-\dots-(k-1)}(x)}{1 - x F_{12-3-\dots-(k-1)}(x)}.$$

Since $F_{12-3-\dots-k}(x) = R_k(x)$ (see 2.3) and $R_k(x)(1 - x R_{k-1}(x)) = 1$, we get $G_{12-3-\dots-k}(x) = x R_k^2(x) G_{12-3-\dots-(k-1)}(x)$, which together with $G_{12}(x) = x^2 R_2^3$ (see 3.3) gives first part of the theorem. $\square$

More generally we can get a formula for $G_{12\dots d-(d+1)-\dots-k}(x)$ as follows.

Theorem 3.5. Let $k \geq d \geq 1$;

$$G_{12\dots d-(d+1)-(d+2)-\dots-k}(x) = \frac{U_d^2\left(\frac{1}{2\sqrt{x}}\right)}{U_k^2\left(\frac{1}{2\sqrt{x}}\right)} G_{[d]}(x).$$

Proof. By use Theorem 3.1 with induction on k (see Corollary 3.4) it follows that

$$G_{12\dots d-(d+1)-\dots-k}(x) = G_{12\dots d}(x) \prod_{j=d+1}^k (x R_j^2(x)).$$

$\square$

Example 3.6. Let $H(x)$ be the generating function for number of all 1-3-2-avoiding permutations such that avoiding 21-3 and containing 21 exactly once; so it is easy to see $H(x) = x^2 F_{21}(x)(F_{21}(x) - 1)$. On the other hand, by Theorem 3.1 we get that $G_{21-3}(x) = xH(x)F_{21}(x) + xF_{21}(x)G_{21-3}(x)$. Hence

$$G_{21-3}(x) = \frac{1}{U_3^2\left(\frac{1}{2\sqrt{x}}\right)}.$$

This example can be generalized by use Theorem 3.1 as following.

Theorem 3.7. For any $k \geq 3$,

$$G_{21-3-\dots-k}(x) = \frac{1}{U_k^2\left(\frac{1}{2\sqrt{x}}\right)}.$$

Proof. By Theorem 3.1 we get that for $k \geq 4$

$$G_{21-3-\dots-k}(x) = \frac{xG_{21-3-\dots-(k-1)}(x)F_{21-3-\dots-k}(x)}{1 - xF_{21-3-\dots-(k-1)}(x)},$$

and by use Theorem 2.3 we get that

$$G_{21-3-\dots-k}(x) = \frac{xR_k(x)}{1 - xR_{k-1}(x)}G_{21-3-\dots-(k-1)}(x),$$

hence, by use the identity $R_k(x)(1 - xR_{k-1}(x)) = 1$ we have that

$$G_{21-3-\dots-k}(x) = xR_k^2(x)G_{21-3-\dots-(k-1)}(x).$$

By use induction on k and Example 3.6 the theorem holds. $\square$

One can try to obtain results similar to Theorems 3.4, 3.5 and 3.7, but expressions involved become extremely cumbersome. So we just consider a simplest cases.

4. CONTAINING 1-3-2 EXACTLY ONCE AND AVOIDING ANOTHER GENERALIZED PATTERN

Denote by $H_\tau(x)$ the generating function for the number of permutations in S_n avoiding a generalized pattern τ and containing 1-3-2 exactly once. We start from the following result, which is analog of result obtained in [MV1].

Theorem 4.1. For any $k \geq 2$,

$$H_{12-3-\dots-k}(x) = \frac{x}{U_k^2\left(\frac{1}{2\sqrt{x}}\right)} \sum_{j=1}^{k-2} U_j^2\left(\frac{1}{2\sqrt{x}}\right)$$

Proof. Let $\alpha \in S_n$ contain 132 exactly once. There are two possibilities: either the only occurrence of 132 in α contains n , or does not contain it. In the first case we get immediately that any entry of α to the right of n is less than any entry of α to the left of n , since otherwise one gets an occurrence of 132 involving n . In the second case, let $i n j$ be the occurrence of 132 in α . First, we have $j = i + 1$, since otherwise either $i + 1 n j$ or $i n i + 1$ would be a second occurrence of 132 in α . Next, i immediately precedes n in α , since if l lies between i and n , then either $l n j$ or $i l j$ would be a second occurrence of 132 in α . Finally, any entry to the right of j is less than any entry between n and j , which in its turn, is less than any entry to the left of i (the proof is similar to the analysis of the first case). Therefore, we have exactly three possibilities for the structure of α :

(i) there exists t such that $\alpha = (\alpha', n, \alpha'')$, where α' is a permutation of $n-1, n-2, \dots, n-t+1$ containing 132 exactly once, and α'' is a permutation of $1, 2, \dots, n-t$ avoiding 132;

(ii) there exists t such that $\alpha = (\alpha', n, \alpha'')$, where α' is a permutation of $n-1, n-2, \dots, n-t+1$ avoiding 132, and α'' is a permutation of $1, 2, \dots, n-t$ containing 132 exactly once;

(iii) there exist t, u such that $\alpha = (\alpha', n-t+1, n, \alpha'', n-t+2, \alpha''')$, where α' is a permutation of $n-1, n-2, \dots, n-t+3$ avoiding 132, α'' is a permutation of $n-t, n-t-1, \dots, n-u+1$ avoiding 132, and α''' is a permutation of $1, 2, \dots, n-u$ avoiding 132.

We are ready to write an equation for $H_{12-3-\dots-k}(x)$ with $k \geq 3$. The contribution of the first structure above is $xH_{12-3-\dots-(k-1)}(x)F_{12-3-\dots-k}(x)$. The contribution of the second possible structure is $x\bar{F}_{12-3-\dots-(k-1)}(x)H_{12-3-\dots-k}(x)$. Finally, the contribution of the third possible structure is $x^3\bar{F}_{12-3-\dots-(k-1)}^2(x)\bar{F}_{12-3-\dots-k}(x)$.

Solving the obtained linear equation and using Theorem 2.3, $H_{12}(x) = 0$ and well known identities involving Chebyshev polynomials (see e.g. [MV3]), we get the desired expression for $H_{12-3-\dots-k}(x)$, $k \geq 3$. $\square$

Example 4.2. Theorem 4.1 yields $H_{12-3}(x) = \frac{1}{U_3^2\left(\frac{1}{2\sqrt{x}}\right)}$. Similarly by the structure of permutations containing exactly 1-3-2 (see proof of Theorem 4.1) we obtain that $H_{21-3}(x) = \frac{1-x}{U_3^2\left(\frac{1}{2\sqrt{x}}\right)}$.

By this example with use the proof of Theorem 4.1, we can be extended this example to generalized pattern $21-3-\dots-k$

Theorem 4.3. For all $k \geq 3$,

$$H_{21-3-\dots-k}(x) = \frac{x}{U_k^2\left(\frac{1}{2\sqrt{x}}\right)} \left(\sum_{j=1}^{k-2} U_j^2\left(\frac{1}{2\sqrt{x}}\right) - 1 \right)$$

One can try to obtain results similar to Theorems 4.1 and 4.3, but expressions involved become extremely cumbersome. So we just consider a two simplest cases.

5. CONTAINING 1-3-2 AND ANOTHER GENERALIZED PATTERN EXACTLY ONCE

Denote by $\Phi_\tau(x)$ the generating function for the number of permutations in $\mathcal{S}_n$ containing both 1-3-2 and a generalized pattern τ exactly once. We start from the following result.

Theorem 5.1. For any $k \geq 2$,

$$\Phi_{12-3-\dots-k}(x) = \frac{1}{U_2(t)U_k^2(t)} \left[1 + \sum_{i=2}^{k-1} \frac{2\sqrt{x}}{U_i(t)U_{i+1}(t)} \left(\sum_{j=1}^i U_j^2(t) - 1 \right) \right], \quad t = \frac{1}{2\sqrt{x}}.$$

Proof. Let $\beta^k = \tau-3-\dots-k$. The three possible structures of permutations containing 1-3-2 exactly once are described in the proof of Theorem 4.1. Let us find the recursion for $\Phi_{\beta^k}(x)$. It is easy to see that the contribution of the first structure equals

$$x\Phi_{\beta^{k-1}}(x)F_{\beta^k}(x) + xH_{\beta^{k-1}}(x)G_{\beta^k}(x),$$

the contribution of the second structure equals

$$xG_{\beta^{k-1}}(x)H_{\beta^k}(x) + xF_{\beta^{k-1}}(x)\Phi_{\beta^k}(x),$$

while the contribution of the third structure equals

$$2x^3G_{\beta^{k-1}}(x)F_{\beta^{k-1}}(x)F_{\beta^k}(x) + x^3F_{\beta^{k-1}}^2(x)G_{\beta^k}(x).$$

Solving the obtained recursion with the initial condition $\Phi_{12}(x) = \frac{x^3}{(1-x)^3}$ (which easy to compute by definitions), and using Theorems 4.1, 2.3, and 3.4, we get the desired result. $\square$

Another interesting results is $\Phi_{21-3-\dots-k}(x)$. Similarly to proof of the above theorem with initial condition $\Phi_{21-3}(x) = \frac{2x^4(1-x)^2}{(1-2x)^3}$ (as a remark $\Phi_{21}(x) = \frac{x^3}{(1-x)^2}$), and using Theorems 2.3, 3.7 and 4.3, we obtain the following.

Theorem 5.2.

$$\Phi_{21-3-\dots-k}(x) = \frac{1}{4t^3U_k^2(t)} \left[\frac{U_2^2(t)}{U_3(t)} + \sum_{i=3}^{k-1} \frac{1}{U_i(t)U_{i+1}(t)} \left(\sum_{j=1}^i U_j^2(t) - 2 \right) \right], \quad t = \frac{1}{2\sqrt{x}}.$$

Remark 5.3. In the above sections, we calculated either $F_\tau(x)$, or $G_\tau(x)$, or $H_\tau(x)$, or $\Phi_\tau(x)$ for given τ generalized patterns of k -letters. We note that there are others interesting cases of τ which we can calculate these functions in the same methods. So we just consider a simplest and interesting cases.

REFERENCES

- [B] M. BÖNA, The permutation classes equinumerous to the smooth class, *Electron. J. Combin.* **5** (1998), #R31
- [BS] E. BABSON AND E. STEINGRIMSSON, Generalized permutation patterns and a classification of the Mahonian statistics, *Séminaire Lotharingien de Combinatoire*, B44b:18pp, (2000).
- [BDPP] E. BARCUCCI, A. DEL LUNGO, E. PERGOLA, AND R. PINZANI, Directed animals, forests and permutations, *Disc. Math.* **204** (1999) 41–71.
- [C] A. CLAEISSON, Generalised pattern avoidance, *European Journal of Combinatorics*, **22** (2001) 961–973.
- [CM] A. CLAEISSON AND T. MANSOUR, Permutations avoiding a pair of generalized patterns of length three with exactly one dash, preprint CO/0107044.
- [CW] T. CHOW AND J. WEST Forbidden subsequences and Chebyshev polynomials *Discr. Math.* **204** (1999) 119–128.
- [Ki] S. KITAEV, Multi-Avoidance of generalised patterns, to appear.
- [Kn] D.E. KNUTH, *The Art of Computer Programming*, 2nd ed. Addison Wesley, Reading, MA (1973).
- [Kr] C. KRATTENTHALER, Permutations with restricted patterns and Dyck paths (2000), preprint CO/0002200
- [K] D. KREMER, Permutations with forbidden subsequences and a generalized Schröder number, *Disc. Math.* **218** (2000), 121–130.
- [MV1] T. MANSOUR AND A. VAINSHTEIN, Restricted permutations, continued fractions, and Chebyshev polynomials, *The Electronic Journal of Combinatorics* **7** (2000), #R17.
- [MV2] T. MANSOUR AND A. VAINSHTEIN, Restricted 132-avoiding permutations, *Adv. Appl. Math.* **126** (2001), 258–269.
- [MV3] T. MANSOUR AND A. VAINSHTEIN, Layered restrictions and Chebychev polynomials (2000), *Annals of Combinatorics*, to appear (2001), preprint CO/0008173.
- [SS] R. SIMION AND F.W. SCHMIDT, Restricted permutations, *European Journal of Combinatorics* **6** (1985) 383–406.
- [R] A. ROBERTSON, Permutations containing and avoiding 123 and 132 patterns, *Disc. Math. and Theo. Comp. Sci.* **3** (1999), 151–154.

- [RWZ] A. ROBERTSON, H. WILF, AND D. ZEILBERGER, Permutation patterns and continuous fractions, *Electron. J. Combin.* **6** (1999), #R38.
- [W] J. WEST, Generating trees and forbidden subsequences, *Disc. Math.* **157** (1996), 363–372.